

BICONFORMAL CHANGES OF METRIC AND PSEUDO-HARMONIC MORPHISMS

RADU SLOBODEANU

ABSTRACT. Pseudo-harmonic morphisms give rise on the domain space to a distribution which admits an almost complex structure compatible with the given Riemannian metric. We shall show that this property, together with the harmonicity, are preserved by a biconformal change of the domain metric. The special case of the pseudo-horizontally homothetic harmonic morphisms is also treated.

1. INTRODUCTION

Pseudo-harmonic morphisms are a special class of harmonic maps into a Hermitian manifold with the additional property called *Pseudo Horizontal Weak Conformality* (PHWC), cf. [6], [7]. This property states that the following relation holds:

$$(1.1) \quad [d\varphi \circ d\varphi^*, J] = 0,$$

where $\varphi : (M, g) \longrightarrow (N, J, h)$ and $d\varphi^*$ denotes the adjoint map.

The geometric meaning of (1.1) will become more transparent if we remark that the differential of a submersion φ from a Riemannian manifold (M, g) into a Hermitian manifold (N, J, h) induces an almost complex structure on the horizontal bundle, defined by $J_{\mathcal{H}} = d\varphi|_{\mathcal{H}}^{-1} \circ J \circ d\varphi|_{\mathcal{H}}$. One can prove that PHWC condition is equivalent to the compatibility of $J_{\mathcal{H}}$ with domain metric g .

Moreover, if the codomain is Kähler, pseudo-harmonic morphisms have a nice description similar to harmonic morphisms. According to [7], pseudo-harmonic morphisms are those maps that pull back local complex-valued holomorphic functions on N to local harmonic functions on M .

Then, the analogue of horizontal homothety in this context was introduced by M.A. Aprodu, M. Aprodu and V. Brinzanescu in [3]. A *Pseudo-Horizontally Homothetic* (PHH) map is a PHWC map φ which satisfies:

$$(1.2) \quad [d\varphi \circ \nabla_X^M \circ d\varphi^*, J] = 0, \forall X \in \Gamma(\mathcal{H})$$

In turn, this condition means that $J_{\mathcal{H}}$ is parallel (with respect to $\nabla^{\mathcal{H}}$) in horizontal directions, so satisfies a *transversal Kähler* condition. Any PHH harmonic submersion onto a Kähler manifold exhibits a particularly nice geometric property: it pulls back complex submanifolds into minimal submanifolds, cf. [3]. Moreover, in [1] it is shown that PHH harmonic submersions are (weakly) stable (in particular such maps minimise the energy-functional in their homotopy class). Further properties and examples of PHH harmonic submersions can be found in [2], [4].

1991 *Mathematics Subject Classification.* 53C12, 58E20.

Key words and phrases. Riemannian manifolds, pseudo-harmonic morphisms, distribution.

In this paper we shall explore the invariance of the above two geometric conditions (PHWC and PHH) together with harmonicity under certain changes of metric. These results are particularly interesting in finding new examples of pseudo-harmonic morphisms and also they point out a nice analogy with the case of harmonic morphisms.

2. CHANGES OF METRIC WHICH PRESERVE PSEUDO-HARMONIC MORPHISMS

According to [8], for a PHWC map $\varphi : (M^m, g) \longrightarrow (N^{2n}, J, h)$ from a Riemannian manifold to a Kähler one, the tension field is given by:

$$\tau(\varphi) = -d\varphi(F^\varphi \delta F^\varphi),$$

where F^φ is the f -structure on M , naturally induced by the PHWC map φ (in fact, F^φ extends $J_{\mathcal{H}}$ with zero on $\mathcal{V}$) and $\delta F^\varphi = \text{trace } \nabla F^\varphi$ is the divergence of F^φ .

If we consider an adapted frame $\{e_i, F^\varphi e_i, e_\alpha\}$ (i.e. an orthonormal frame such that $e_\alpha \in \text{Ker } F^\varphi$), then the local form of the above formula is:

$$(2.1) \quad \tau(\varphi) = -d\varphi \left(\sum_{i=1}^n F^\varphi [(\nabla_{e_i} F^\varphi)(e_i) + (\nabla_{F^\varphi e_i} F^\varphi)(F^\varphi e_i)] + (m - 2n)\mu^\mathcal{V} \right).$$

Definition 2.1. Let (M, g) be a Riemannian manifold endowed with two complementary distributions $\mathcal{V}$ and $\mathcal{H}$ which induce a natural decomposition of the metric: $g = g^\mathcal{V} + g^\mathcal{H}$. Then one call *biconformal change of metric* g , the association of a new metric on M defined by:

$$\bar{g} = \sigma^{-2} g^\mathcal{H} + \rho^{-2} g^\mathcal{V},$$

where $\sigma, \rho : M \longrightarrow (0, \infty)$ are smooth functions.

The next result for PHWC submersions is similar to that one for horizontally conformal submersions (see [5], Lemma 4.6.6, p. 126).

Proposition 2.1. *Let $\varphi : (M^m, g) \longrightarrow (N^{2n}, J, h)$ be a PHWC submersion. With respect to the vertical and horizontal distributions, one take a biconformal change of the metric g , denoted by $\bar{g}$. Then, regarded as a map from $(M, \bar{g})$ to (N^{2n}, J, h) , φ has tension field:*

$$(2.2) \quad \bar{\tau}(\varphi) = \sigma^2 [\tau(\varphi) + d\varphi(\text{grad}(\ln \rho^{2n-m} \sigma^{2-2n}))]$$

Proof. Consider an adapted frame $\{e_i, F^\varphi e_i, e_\alpha\}$ for (M, g) . Then $\{\sigma e_i, \sigma F^\varphi e_i, \rho e_\alpha\}$ will be an adapted frame with respect to the perturbed metric $\bar{g}$. Let f_i denote e_i when $i = \overline{1, n}$ and $F^\varphi e_{i-n}$ when $i = \overline{n+1, 2n}$. Then, using Koszul formula, one obtain for any $X, Y \in \Gamma(\mathcal{H})$:

$$(2.3) \quad \mathcal{H}(\bar{\nabla}_X Y) = \mathcal{H}(\nabla_X Y) + \sum_{i=1}^n [-X(\ln \sigma)g(Y, f_i) - Y(\ln \sigma)g(X, f_i) + f_i(\ln \sigma)g(X, Y)] f_i.$$

In the same way, but for $V \in \Gamma(\mathcal{V})$ one get:

$$(2.4) \quad \mathcal{H}(\bar{\nabla}_V V) = \frac{\sigma^2}{2} \left[2\rho^{-2} \mathcal{H}(\nabla_V V) + \sum_{i=1}^n [-f_i(\rho^{-2})g(V, V)] f_i \right].$$

Now, applying (2.4) for $V = \rho e_\alpha, \alpha = \overline{2n+1, m}$ and summing, one get the following result which has the same form as in the harmonic morphism case (see [5], Remark 4.6.8, p. 126):

Lemma 2.1. *At a biconformal change of the metric, the mean curvature of the fibers of a pseudo-harmonic morphism changes in this way:*

$$(2.5) \quad \bar{\mu}^\mathcal{V} = \sigma^2 [\mu^\mathcal{V} + \mathcal{H}(\text{grad}(\ln \rho))]$$

Now, in order to get the formula stated by the theorem, one has to see how changes the first term in equation (2.1), that is:

$$\begin{aligned} F^\varphi \text{div}_\mathcal{H} F^\varphi &= F^\varphi [(\nabla_{e_i} F^\varphi)(e_i) + (\nabla_{F^\varphi e_i} F^\varphi)(F^\varphi e_i)] = \\ &\mathcal{H}(\nabla_{e_i} e_i) + \mathcal{H}(\nabla_{F^\varphi e_i} F^\varphi e_i) + F^\varphi(\nabla_{e_i} F^\varphi e_i) - F^\varphi(\nabla_{F^\varphi e_i} e_i). \end{aligned}$$

So we have to apply formula (2.3) in four particular cases:

$$\begin{aligned} \mathcal{H}(\bar{\nabla}_{\sigma e_i} \sigma e_i) &= \sigma^2 \mathcal{H}(\nabla_{e_i} e_i) - \sigma^2 e_i(\ln \sigma) e_i + \sigma^2 \text{grad}_\mathcal{H}(\ln \sigma), \\ \mathcal{H}(\bar{\nabla}_{\sigma F^\varphi e_i} \sigma F^\varphi e_i) &= \sigma^2 \mathcal{H}(\nabla_{F^\varphi e_i} F^\varphi e_i) - \sigma^2 F^\varphi e_i(\ln \sigma) F^\varphi e_i + \sigma^2 \text{grad}_\mathcal{H}(\ln \sigma), \\ \mathcal{H}(\bar{\nabla}_{\sigma e_i} \sigma F^\varphi e_i) &= \sigma^2 \mathcal{H}(\nabla_{e_i} F^\varphi e_i) - \sigma^2 F^\varphi e_i(\ln \sigma) e_i, \\ \mathcal{H}(\bar{\nabla}_{\sigma F^\varphi e_i} \sigma e_i) &= \sigma^2 \mathcal{H}(\nabla_{F^\varphi e_i} e_i) - \sigma^2 e_i(\ln \sigma) F^\varphi e_i. \end{aligned}$$

Now, summing these four terms as they appear in $F^\varphi \bar{\text{div}}_\mathcal{H} F^\varphi$ we get:

$$(2.6) \quad F^\varphi \bar{\text{div}}_\mathcal{H} F^\varphi = \sigma^2 (F^\varphi \text{div}_\mathcal{H} F^\varphi + (2n - 2) \text{grad}(\ln \sigma))$$

Now, from (2.5) and (2.6), the stated result follows. $\square$

The following result is analogous with the one proved in [9].

Corollary 2.1. *Let $\varphi : (M^m, g) \longrightarrow (N^{2n}, J, h)$ be a PHWC submersion (with $m > 2n$). For any smooth function $\sigma : M \longrightarrow (0, \infty)$, set*

$$(2.7) \quad g_\sigma = \sigma^{-2} g^\mathcal{H} + \sigma^{\frac{4n-4}{m-2n}} g^\mathcal{V}.$$

Then φ is a pseudo-harmonic morphism with respect to g if and only if is a a pseudo-harmonic morphism with respect to g_σ .

Now, suppose that φ is a pseudo-horizontally homothetic map. The condition (1.2) translates simply into $d\varphi((\nabla_X F^\varphi)Y) = 0, \forall X, Y \in \Gamma(\mathcal{H})$, so:

$$[(\nabla_X F^\varphi)Y]^\mathcal{H} = 0, \forall X, Y \in \Gamma(\mathcal{H}).$$

In particular, $F^\varphi \text{div}_\mathcal{H} F^\varphi = 0$, so the formula (2.1) reduces to:

$$\tau(\varphi) = -(m - 2n) d\varphi(\mu^\mathcal{V}).$$

From the above relation, one can see that a pseudo-horizontally homothetic map is harmonic if and only if the fibers are minimal (as it was proved in [3]).

We want to see under which changes of metric of the type (2.7) the PHH harmonic morphisms are invariant.

Corollary 2.2. *Let $\varphi : (M^m, g) \longrightarrow (N^{2n}, J, h)$ be a PHH harmonic morphism (with $m > 2n$). Then φ is a PHH harmonic morphism with respect to g_σ if and only if σ is constant. In particular, for any any strictly positive constant c , set*

$$g_c = c^{-2} g^\mathcal{H} + c^{\frac{4n-4}{m-2n}} g^\mathcal{V}.$$

Then a PHH submersion φ is a PHH harmonic morphism with respect to g if and only if is a a PHH harmonic morphism with respect to g_c .

Proof. In particular, φ is a PHWC submersion. Using (2.3), for any X tangent to $\mathcal{H}$ we derive:

$$\begin{aligned} \mathcal{H}((\overline{\nabla}_X F^\varphi)Y) &= \mathcal{H}((\nabla_X F^\varphi)Y) + \text{grad}_{\mathcal{H}}(\ln\sigma)g(X, F^\varphi)Y + \\ &g(X, Y) \sum_{i=1}^n [F^\varphi e_i(\ln\sigma)e_i + e_i(\ln\sigma)F^\varphi e_i] - \\ &F^\varphi Y(\ln\sigma)X + Y(\ln\sigma)F^\varphi X. \end{aligned}$$

As φ is a PHH harmonic morphism with respect to g , if σ is constant, then it is obvious from the above formula that φ is PHH harmonic morphism with respect to g_σ , too.

Conversely, if φ is PHH harmonic morphism with respect to g_σ , then $F^\varphi \overline{\text{div}}_{\mathcal{H}} F^\varphi = 0$. But the hypothesis implies also $F^\varphi \text{div}_{\mathcal{H}} F^\varphi = 0$ and the relation (2.6) tells us precisely that σ must be a constant. $\square$

REFERENCES

- [1] M.A. Aprodu, *PHH harmonic submersions are energy-minimizing*, Preprint.
- [2] M.A. Aprodu, M. Aprodu, *Implicitly defined harmonic PHH submersions*, Manuscripta Math. **100** (1999), 103-121.
- [3] M.A. Aprodu, M. Aprodu, V. Brinzanescu, *A class of harmonic submersions and minimal submanifolds*, Int. J. Math. **11**(9) (2000), 1177-1191.
- [4] V. Brinzanescu, *Pseudo-harmonic morphisms; applications and examples*, The Fifth International Workshop on Differential Geometry and Its Application, Sept. 18-22, 2001, Timisoara, Romania, in *Analele Univ.Timisoara*, Vol. **XXXIX** (2001), 111-121.
- [5] P. Baird, J. C. Wood, *Harmonic Morphisms Between Riemannian Manifolds*, Clarendon Press - Oxford (2003).
- [6] J. Chen, *Structures of certain harmonic maps into Kähler manifolds*, Int. J. Math. **8** (1997), 573-581.
- [7] E. Loubeau, *Pseudo harmonic morphisms*, Int. J. of Math. **8** (1997), 943-957.
- [8] E. Loubeau, X. Mo, *Pseudo horizontally weakly conformal maps from Riemannian Manifolds into Kähler Manifolds*, Beiträge zur Alg. und Geom. (to appear).
- [9] X. Mo, *On the geometry of horizontally homothetic maps and harmonic morphisms*, Adv. Math. Beijing **23** (1994), 282-283.

FACULTY OF PHYSICS, BUCHAREST UNIVERSITY, ROMANIA
E-mail address: `slobir@k.ro`